

# Introgression of *Aegilops speltoides* segments in *Triticum aestivum* and the effect of the gametocidal genes

## Julie King<sup>1,\*</sup>, Surbhi Grewal<sup>1</sup>, Cai-yun Yang<sup>1</sup>, Stella Hubbart Edwards<sup>1</sup>, Duncan Scholefield<sup>1</sup>, Stephen Ashling<sup>1</sup>, John A. Harper<sup>2</sup>, Alexandra M. Allen<sup>3</sup>, Keith J. Edwards<sup>3</sup>, Amanda J. Burridge<sup>3</sup> and Ian P. King<sup>1</sup>

<sup>1</sup>Division of Plant and Cop Sciences, University of Nottingham, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, UK, <sup>2</sup>The Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Plas Gogerddan, Aberystwyth SY23 3EB, UK and <sup>3</sup>School of Biological Sciences, University of Bristol, Bristol BS8 1TQ, UK \*For correspondence. E-mail julie.king@nottingham.ac.uk

Received: 25 May 2017 Returned for revision: 28 August 2017 Editorial decision: 8 September 2017 Accepted: 13 October 2017 Published electronically: 5 December 2017

• **Background and Aims** Bread wheat (*Triticum aestivum*) has been through a severe genetic bottleneck as a result of its evolution and domestication. It is therefore essential that new sources of genetic variation are generated and utilized. This study aimed to generate genome-wide introgressed segments from *Aegilops speltoides*. Introgressions generated from this research will be made available for phenotypic analysis.

• **Methods** *Aegilops speltoides* was crossed as the male parent to *T. aestivum* 'Paragon'. The interspecific hybrids were then backcrossed to Paragon. Introgressions were detected and characterized using the Affymetrix Axiom Array and genomic *in situ* hybridization (GISH).

Key Results Recombination in the gametes of the F<sub>1</sub> hybrids was at a level where it was possible to generate a genetic linkage map of Ae. speltoides. This was used to identify 294 wheat/Ae. speltoides introgressions. Introgressions from all seven linkage groups of Ae. speltoides were found, including both large and small segments. Comparative analysis showed that overall macro-synteny is conserved between Ae. speltoides and T. aestivum, but that Ae. speltoides does not contain the 4A/5A/7B translocations present in wheat. Aegilops speltoides has been reported to carry gametocidal genes, i.e. genes that ensure their transmission through the gametes to the next generation. Transmission rates of the seven Ae. speltoides linkage groups introgressed into wheat varied. A 100 % transmission rate of linkage group 2 demonstrates the presence of the gametocidal genes on this chromosome.
Conclusions A high level of recombination occurs between the chromosomes of wheat and Ae. speltoides, leading to the gametocidal genes, all germplasm developed will always contain a segment from Ae. speltoides linkage group 2S, in addition to an introgression from any other linkage group.

Key words: *Aegilops speltoides, Triticum aestivum*, introgression, recombination, comparative synteny, gametocidal, genetic linkage mapping, GISH, chromosome transmission

## INTRODUCTION

Bread wheat (*Triticum aestivum*), one of the world's leading sources of food, is an allopolyploid (6x = AABBDD = 42) composed of the genomes of three different species. The A genome is derived from *Triticum urartu* (Dvorak *et al.*, 1993), the B from *Aegilops speltoides* or a closely related species (Sarkar and Stebbins, 1956; Riley *et al.*, 1958; Dvorak and Zhang, 1990; Maestra and Naranjo, 1998; Marcussen *et al.*, 2014) and the D genome from *Aegilops tauschii* (McFadden and Sears, 1946). The final hybridization event between tetraploid wheat (AABB) and *Ae. tauschii* (DD) is thought to have occurred only once or twice, ~8000–10 000 years ago. As a result, wheat went through a significant genetic bottleneck. Thus, the significant yield gains achieved by wheat breeders to date have been via the exploitation of genetic variation that has arisen via gene mutation over the last 8000–10 000 years and rare outcrossing events with tetraploid wheat. In many parts of the world wheat yields are now starting to plateau and this is thought to be a direct result of a lack of genetic variation compounded by the changing environment at a time of increasing demand for food due to the increasing global population (Charmet, 2011; Grassini *et al.*, 2013; Ray *et al.*, 2013). It is therefore essential that new sources of genetic variation be found that will enable breeders to generate the next generation of high-yielding environmentally adapted wheat varieties (Tanksley and McCouch, 1997; Haussmann *et al.*, 2004; Tester and Langridge, 2010; McCouch *et al.*, 2013; Warschefsky *et al.*, 2014; Brozynska *et al.*, 2015; Zhang *et al.*, 2017)

Unlike wheat, its progenitor species and wild relatives provide a vast and largely untapped source of genetic variation for most, if not all, traits of agronomic importance. Considerable efforts have been made to exploit the genetic variation within

(http://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and

reproduction in any medium, provided the original work is properly cited.

<sup>©</sup> The Author(s) 2017. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License

the progenitors and wild relatives, with some noticeable successes, for example the transfer of a segment of Aegilops *umbellulata* to wheat conferring resistance to leaf rust (Sears, 1955), the transfer of a segment from Aegilops ventricosa carrying resistance to eyespot (Maia, 1967; Doussinault et al., 1983) and its subsequent release in the variety Rendevouz, and the registration of 'MACE', a hard red winter wheat carrying a segment from Thinopyrum intermedium conferring resistance to wheat streak mosaic virus (Graybosch et al., 2009). Progress has been severely hindered, however, due to an inability to quickly and accurately identify and characterize interspecific transfers to wheat. However, with the advent of new molecular technologies coupled with specific crossing strategies, we can now systematically exploit the genetic variation available in the progenitors of wheat and its wild relatives. (e.g. Tiwari et al., 2014, 2015; King et al., 2016, 2017).

The transfer of genetic variation to wheat from its progenitors/wild relatives occurs via recombination between the chromosomes of wheat and those of its wild relative at meiosis in  $F_1$  hybrids or in their derivatives, e.g. addition and substitution lines (King *et al.*, 2016, 2017). This results in the generation of interspecific recombinant chromosomes. Once identified, interspecific recombinant chromosomes are then recurrently backcrossed with wheat. The ultimate objective is to generate lines of wheat carrying a small chromosome segment from a progenitor/wild relative with a target gene but few, if any, deleterious genes.

One of the wild relatives of wheat that is of particular interest to researchers is *Ae. speltoides*, a rich source of genetic variation for resistance to a range of diseases of importance in wheat (Jia *et al.*, 1996; Mago *et al.*, 2009; Klindworth *et al.*, 2012; Deek and Distelfeld, 2014). *Aegilops speltoides*, or an extinct close relative, has been proposed as the donor of the B genome of wheat (Sarkar and Stebbins, 1956; Dvorak and Zhang, 1990; Friebe and Gill, 1996; Tsunewaki, 1996; Dvorak and Akhunov, 2005; Marcussen *et al.*, 2014). *Aegilops speltoides* is a facultative outbreeder, but can be readily inbred in the glasshouse.

Unlike most of the wild relatives of wheat, *Ae. speltoides* possesses the ability to suppress the wheat *Ph1* locus (Dvorak, 1972; Dvorak *et al.*, 2006). Recombination is normally restricted to homologous chromosomes from the same genome, i.e. although the A, B and D genomes of wheat and chromosomes from wild relatives are related, recombination can only occur within genomes in the presence of *Ph1*. However, if *Ph1* is removed, related chromosomes (homoeologues) from different genomes can recombine at meiosis. *Aegilops speltoides* possesses genes on chromosomes 3S (*Su1-Ph1*) and 7S (*Su2-Ph1*) (Dvorak *et al.*, 2006), with the ability to suppress *Ph1*. Thus, in  $F_1$  hybrids between wheat and *Ae. speltoides Su1-Ph1* or *Su2-Ph1* enables intergenomic homoeologous recombination to occur during meiosis.

Aegilops speltoides also carries gametocidal genes (Gc) (Tsujimoto and Tsunewaki, 1984, 1988; Ogihara *et al.*, 1994), which are preferentially transmitted to the next generation. Individuals heterozygous for Gc gene(s) produce two types of gametes. Both male and female gametes that lack the Gc genes are generally not viable due to chromosome breakage (Finch *et al.*, 1984). Gametes that carry the Gc genes, however, behave normally and thus it is only these gametes that are transmitted to the next generation. Aegilops speltoides carries two alleles of the gametocidal genes. Gc1a and Gc1b, both located on

chromosome 2S, are transmitted at a very high frequency to the next generation and give rise to different morphological aberrations. Gc1a causes endosperm degeneration and chromosome aberrations while Gc1b results in the production of seeds that lack shoot primordia (Tsujimoto and Tsunewaki, 1988). The limitations of utilizing species with gametocidal genes for wild relative introgression are: (1) the fertility of hybrid/crossed grain is reduced by at least 50 %; and (2) all introgressions from wild relatives with Gc genes will always carry them and any other genes closely linked to them. Hence it is very difficult, if not impossible, to transfer genes from other regions of the genome in the absence of the Gc genes.

In this paper we investigate (1) the ability of the *Ae. spel-toides* suppressors of the *Ph1* locus located on chromosomes 3S and 7S to induce recombination between the chromosomes of wheat and those of *Ae. speltoides*, and (2) the frequency of transmission of the *Ae. speltoides* gametocidal chromosomes through the gametes in  $F_1$  hybrids and their derivatives.

#### MATERIALS AND METHODS

## Generation of introgressions

The method used to induce introgressions between wheat and *Ae. speltoides* is the same as that described by King *et al.* (2017). In summary, hexaploid wheat ('Paragon', obtained from the Germplasm Resources Unit at the JIC; code W10074) was pollinated with *Ae. speltoides* (accession 2140066 obtained from the Germplasm Resources Unit at the JIC) to produce  $F_1$ interspecific hybrids. These hybrids were then grown to maturity and backcrossed as the female with the wheat parent to generate BC<sub>1</sub> populations. The BC<sub>1</sub> individuals and their resulting progenies were then recurrently pollinated to produce BC<sub>2</sub>, BC<sub>3</sub> etc. populations(Fig. 1).

As a result of the presence of the suppressors of the *Ph1* locus, homoeologous recombination was expected to occur at meiosis in the  $F_1$  wheat/*Ae. speltoides* hybrids, leading to the production of interspecific recombinant chromosomes/ introgressions. Subsequent recurrent backcrossing of any BC<sub>1</sub> progeny to the wheat parent was undertaken to transfer any interspecific recombinant chromosomes generated into a wheat (Paragon) background.

#### Detection of putative wheat/Ae. speltoides introgressions

A 35K Axiom<sup>®</sup> Wheat-Relative Genotyping Array (Affymetrix, Santa Clara, CA) was used to detect the presence of putative wheat/*Ae. speltoides* introgressions in each of the backcross generations (King *et al.*, 2017). In summary, the array is composed of single-nucleotide polymorphisms (SNPs), each showing polymorphism for the ten wild relatives (under study at the Nottingham/ BBSRC Wheat Research Centre), including *Ae. speltoides* (King *et al.*, 2017). All the SNPs incorporated in this array formed part of the Axiom<sup>®</sup> 820K SNP array (Winfield *et al.*, 2016). The dataset for the Axiom<sup>®</sup> 820K SNP array is available from www.cerealsdb. uk.net (Wilkinson *et al.*, 2012, 2016). Table 1 shows the number of putative SNPs between *Ae. speltoides* and each of the wheat genotypes included on the array. The array has been constructed in such a way that up to 384 lines could be screened at one time. Thus, the array facilitated the high-throughput, high-resolution



FIG. 1. Wheat/wild relative introgression strategy.

 TABLE 1. Number of polymorphic SNPs between Ae. speltoides and hexaploid wheat based on the Affymetrix 35K array. A comparison is made between all calls (all markers showing polymorphism between wheat and Ae. speltoides) and Poly High Resolution (PHR) calls, which are codominant and polymorphic between wheat and Ae. speltoides with at least two examples of the minor allele

		Linkage group						Total
	1	2	3	4	5	6	7	_
All calls (% of total)	2770 (12.4)	3992 (17.9)	3358 (15.1)	2798 (12.6)	3646 (16.4)	2530 (11.4)	3164 (14.2)	22 258
PHR calls (% of total)	69 (12.7)	78 (14.3)	82 (15.1)	85 (15.6)	90 (16.5)	46 (8.5)	94 (17.3)	544

screening of wheat/*Ae. speltoides* introgressions. Genotyping was performed as described by King *et al.* (2017) with no modifications (see below).

## Genetic mapping of Ae. speltoides

Genetic mapping was as described by King *et al.* (2017); in summary, DNA was extracted using a CTAB method (Zhang *et al.*, 2013) from individuals of the back-crossed populations,  $BC_1$ ,  $BC_2$ ,  $BC_3$ ,  $BC_4$  and  $BC_5$  derived from the wheat/*Ae. speltoides F*<sub>1</sub> hybrids. These populations were genotyped with the Axiom<sup>®</sup> Wheat-Relative Genotyping Array. Only Poly High Resolution (PHR) SNP markers, which were codominant and polymorphic, with at least two examples of the minor allele were used for genetic mapping (King *et al.*, 2017). The SNP markers that showed (1) heterozygous calls for either parent(s), (2) no polymorphism between the wheat parents and *Ae*. *speltoides*, and/or (3) no calls for either parent(s) were removed using Flapjack<sup>TM</sup> (Milne *et al.*, 2010; v.1.14.09.24). The resulting markers were sorted into linkage groups (Fig. 2) in JoinMap<sup>®</sup> 4.1 (van Ooijen, 2011) with a LOD score of 20 and a recombination frequency threshold of 0.1 using the Haldane mapping function (Haldane, 1919). All markers that did not show any heterozygous call or were unlinked were ignored and only the highest-ranking linkage groups with >30 markers were selected for map construction. These were exported and assigned to chromosomes using information from the Axiom<sup>®</sup> Wheat HD Genotyping Array (Winfield *et al.*, 2016).

## Comparative analysis

Synteny analysis was carried out using sequence information of the markers located on the present map of *Ae. speltoides*. The sequences of the mapped markers were compared using

#### Aegilops speltoides linkage groups



FIG. 2. Genetic linkage map of Ae. speltoides.

BLAST (e-value cut-off of 1e-05) against the wheat genome (http://plants.ensembl.org/Triticum\_aestivum) to obtain the orthologous map positions of the top hits in the A, B and D genomes of wheat. To generate the figures, centimorgan (cM) distances on the linkage groups of the present map of *Ae. speltoides* were scaled up by a factor of 100 000 to match similar base-pair lengths of the chromosomes of the wheat genome. Figure 3 was visualized using Circos (v. 0.67; Krzywinski *et al.*, 2009) in order to observe synteny between *Ae. speltoides* (genetic position in cM) and the wheat genome (physical position in Mb).

#### Cytogenetic analysis

The protocol for genomic *in situ* hybridization (GISH) was as described in Zhang *et al.* (2013), Kato *et al.* (2004) and King *et al.* (2017). In summary, genomic DNAs from young leaves of the three putative diploid progenitors of bread wheat, i.e. *T. urartu* (A genome), *Ae. speltoides* (B genome) and *Ae. tauschii* (D genome), were isolated using extraction buffer (0.1 M Tris–HCl pH 7.5, 0.05 M EDTA pH 8.0, 1.25% SDS). Samples were incubated at 65 °C for 1 h before being placed on ice and mixed with ice-cold 6 M NH<sub>4</sub>C<sub>2</sub>H<sub>3</sub>O for 15 min. The samples were then spun down, the supernatant was mixed with

#### Aegilops speltoides linkage groups



FIG. 2. Continued.

isopropanol to pellet the DNA and the isolated DNA was further purified with phenol/chloroform. The genomic DNAs of *Ae. speltoides* and *T. urartu* were labelled by nick translation with Chroma Tide Alexa Fluor 488-5-dUTP (Invitrogen, Carlsbad, CA; C11397). Genomic DNA of *Ae. tauschii* was labelled with Alexa Fluor 594-5-dUTP (Invitrogen; C11400). Genomic DNAs of *Ae. speltoides* and *T. aestivum* 'Chinese Spring' were fragmented to 300–500 bp in a heat block at 100 °C.

Preparation of chromosome spreads was as described in Kato *et al.* (2004) and King *et al.* (2017). Briefly, roots were excised from germinated seeds, treated with nitrous oxide gas at 10 bar for 2 h, fixed in 90 % acetic acid for 10 min and then washed three times in water on ice. Root tips were dissected and digested in 20  $\mu$ L of 1 % pectolyase Y23 and 2 % cellulase Onozuka R-10 (Yakult Pharmaceutical, Tokyo) solution for 50

min at 37 °C and then washed three times in 70 % ethanol. Root tips were crushed in 70 % ethanol, cells collected by centrifugation at 2.5 g for 1 min, briefly dried and then re-suspended in 30–40  $\mu$ L of 100 % acetic acid prior to being placed on ice. The cell suspension was dropped onto glass slides (6–7  $\mu$ L per slide) in a moist box and dried slowly under cover.

Slides were initially probed using labelled genomic DNA of *Ae. speltoides* (100 ng) and fragmented genomic DNA of Chinese Spring (3000 ng) as blocker (in a ratio of 1: 30 per slide) to detect the *Ae. speltoides* introgressions. The slides were bleached (dipped in 2 × saline–sodium citrate (SSC) to remove the coverslip, transferred to  $4 \times$  SSC for 5 min and air-dried in the light) and re-probed with labelled DNAs of *T. urartu* (100 ng) and *Ae. tauschii* (200 ng) and fragmented DNA of *Ae. speltoides* (3000 ng) as blocker in a ratio of 1: 2:30 per



FIG. 3. (A) SNP characterization of *Ae. speltoides* introgressions in five consecutive generations, i.e.  $BC_1$ ,  $BC_2$ ,  $BC_3$ ,  $BC_4$  and  $BC_5$ . (B) Genomic *in situ* hybridization image of the  $BC_5$  genotype (aneuploid 41-chromosome plant). In the SNP characterization red colour is used to represent the presence of an *Ae. speltoides* introgression and blue colour represents wheat. It should be noted that these diagrams cannot be used to assess which wheat chromosomes the *Ae. speltoides* segments have recombined with. The GISH image shows a metaphase spread of  $BC_5$ -5A probed with labelled genomic DNA of *Ae. speltoides*. Arrows show *Ae. speltoides* introgressions (green).

slide to detect the AABBDD genomes of wheat. In both cases, the hybridization mix was made up to 10  $\mu$ L with 2 × SSC in 1 × Tris/EDTA buffer (TE). Slides were incubated initially at 75 °C for 5 min and then overnight at 55 °C in a closed box

before counterstaining with Vectashield mounting medium with 4',6-diamidino-2-phenylindole,dihydrochloride (DAPI), and analysed using a high-throughput, fully automated Zeiss Axio Imager Z2 upright epifluorescence microscope (Carl Zeiss, Oberkochen, Germany) with filters for DAPI (blue), Alexa Fluor 488 (green) and Alexa Fluor 594 (red). Photographs were taken using a MetaSystems Coolcube 1m CCD camera. Further slide analysis was carried out using Meta Systems ISIS and Metafer software (Metasystems, Altlussheim, Germany). This system enabled the fully automated capture of high- and lowpower fluorescent images of root tip metaphase spreads. Slides with root tip preparations were automatically scanned and the images downloaded for analysis.

#### RESULTS

#### Generation of wheat/Ae. speltoides introgressions

A total of 3890 crosses were made between wheat and *Ae. speltoides* and their derivatives, leading to the generation of 9953 crossed seeds and 2029 self-seeds (Fig. 1). The number of seeds germinated, plants crossed and seed set are shown in Table 2. Every ear produced by the  $F_1$  was crossed with wheat. The  $F_1$ hybrids generated between wheat and *Ae. speltoides* showed the lowest frequency of germination. Only 15 % germinated compared with 100, 73, 70 and 78 % in the BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub> and BC<sub>4</sub> generations. The  $F_1$  hybrids also showed the lowest fertility, with only 29 % of crossed ears producing any seeds compared with crossed ears from the BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub> and BC<sub>5</sub> generations, which showed 58, 72, 83 and 84 % fertility, respectively.

Of the 20  $F_1$  seeds germinated, only three reached maturity and set seeds. Thus all of the wheat/Ae. speltoides introgressions generated in this programme were limited to these three viable  $F_1$  hybrids. Table 2 summarizes the number of seeds germinated and seed set for each of the backcross generations. In each generation, the average seed set per crossed ear was very low, i.e. the lowest average seed set per ear was 0.6 (observed in the  $F_1$  hybrid), while the highest average seed set per crossed ear was only 3.7 (observed in the BC<sub>2</sub>). The fertility of plants that had previously been self-fertilized at least once was considerably higher than that of plants in the backcross programme, although again variability was seen, e.g.  $BC_3F_1$ -16A was male-sterile and therefore produced no self-fertilized seed, while seed numbers from crossed ears ranged from 8 to 19. Some plants were also found to be asynchronous for anther and stigma maturity. For these plants with manual pollination, self-fertilized seeds were produced.

#### Detection of introgressions

Of the SNPs on the Axiom array, 22 258 were polymorphic between *Ae. speltoides* and wheat (Table 1). The Axiom array was used to screen genomic DNA prepared from 536 samples of BC<sub>1</sub> to BC<sub>5</sub> lines between wheat and *Ae. speltoides*. Genotype calls were generated, and the sample call rate (markers working in a particular genotype) ranged from 86.5 to 99.8 %, with an average of 98.8 % for the 536 samples. Affymetrix software classified the scores for each SNP into one of six cluster patterns. However, only those classified as Poly High Resolution (PHR) were used for genetic mapping as these are considered to be optimum quality. Linkage group 7 had the highest number of SNPs (17.3 %), while linkage group 6 had the lowest number (8.5 %).

JoinMap<sup>®</sup> (van Ooijen, 2011) was used to analyse the PHR SNPs and this led to the establishment of seven linkage groups. The genetic map was composed of 544 SNPs and represented the seven chromosomes of *Ae. speltoides*. Linkage groups 1-7 had 69, 78, 82, 85, 90, 46 and 94 SNPs respectively (Fig. 2). The lengths of linkage groups 1-7 were 70.8, 90.3, 98.5, 68, 110.5, 63.6 and 92.9 cM, respectively, with a total length of 594.6 cM and an average chromosome length of 84.9 cM.

## Genomic in situ hybridization

In most cases the number of wheat/*Ae. speltoides* introgressions detected by genomic *in situ* hybridization (GISH) in  $BC_4$  and  $BC_5$  individuals corresponded exactly with the number detected via SNP analysis (Table 3 and Fig. 3). One exception was detected in the GISH analysis of  $BC_5$ -4A. The SNP analysis of this plant showed the only segment present to be a complete 2S chromosome. However, the GISH image showed the presence of a large *Ae. speltoides* 2S segment (one telomere was missing from the chromosome) and a very small *Ae. speltoides* segment recombined with a different wheat chromosome.

#### Syntenic relationship

Figure 4 shows the syntenic relationship between the seven linkage groups of *Ae. speltoides* and the three genomes of wheat, with large ribbons showing significant synteny. Some

 

 TABLE 2. Number of seed produced and germinated in relation to the number of crosses carried out for each generation of the introgression programme for Ae. speltoides into wheat

	Wheat × Ae. speltoides	$F_{1}$	BC <sub>1</sub>	BC <sub>2</sub>	BC <sub>3</sub>	$\mathrm{BC}_4$	Total
Number of seeds sown	NA	20	22	400	187	173	802
Number of seeds that germinated (%)	NA	3 (15)	22 (100)	292 (73)	130 (70)	135 (78)	582
Number of plants that set seed (%)	NA	3 (100)	22 (100)	261 (89)	124 (95)	117 (87)	527
Number of seed/total number of crosses (average number of seed set per crossed ear)	127/26 (4.9)	22/35 (0.6)	509/357 (1.4)	5743/2180 (2.6)	3204/876 (3.7)	348/416 (0.8)	9963/3890
Number of crosses producing seed (%)	18 (69)	10 (29)	207 (58)	1579 (72)	730 (83)	350 (84)	2894
Number of self seed produced	0	0	21	520	753	735	2029

TABLE 3. Number of introgressed segments from Ae. speltoides present in  $BC_4$  and  $BC_5$  plants as detected by SNP genotyping and GISH. The Ae. speltoides chromosomes have been assigned to linkage groups via the comparative analysis of the SNPs with wheat

Plant accession numbers		Number of	Ae. speltoides	
Backcross	Number	Genotyping	GISH	segments
BC <sub>4</sub> BC <sub>5</sub>	45A	3	3	2, 5, 6
	45C	1	1	2
	45D	3	3	2, 5, 6
	1A	3	3	2, 4, 5
	1B	1	1	2
	1C	1	1	2
	1D	2	2	2,5
	2	2	2	2,5
	3	1	1	1
	4A	1	1 + small	2,?
			end	
	5A	2	2	1, 2
	6A	1	1	2
	6B	2	2	1,2
	6C	1	1	2
	7A	2	2	2, 5
	7C	1	1	2
	8A	1	1	2
	8B	1	1	2
	8C	1	1	2
	8D	1	1	2

gene rearrangements are indicated where single markers crossmap to non-collinear positions on wheat chromosomes. The only major disruption in synteny between the two species is that *Ae. speltoides* does not carry the 4/5/7 translocation observed for chromosomes 4A, 5A and 7B of wheat (Liu *et al.*, 1992; Naranjo *et al.*, 1987). These data demonstrate that there is a close syntenic relationship between the A, B and D genomes of wheat and *Ae. speltoides*.

## Preferential transmission

The data obtained from SNP analysis was used to determine the transmission frequency of chromosome segments from each of the seven *Ae. speltoides* linkage groups through the female gametes to the BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub>, BC<sub>4</sub> and BC<sub>5</sub> generations. Chromosome segments from linkage group 2 were observed in all 536 backcross progeny observed (Table 4). The smallest segment carried only three SNP markers, AX94631566, AX95123223 and AX94601746, located near one of the terminal ends of the 2S linkage group (Fig. 2). These three SNP markers were found to be present in all the backcross individuals analysed. In order to determine whether the 2S chromosome was also preferentially transmitted through the male gametes, 41 BC<sub>3</sub> and BC<sub>4</sub> plants were allowed to selffertilize and their progenies were analysed using GISH. The BC<sub>3</sub>/BC<sub>4</sub> plants selected each carried a single copy of a 2S chromosome segment. Thus, these plants would produce two types of male and female gametes during gametogenesis: those that carried 2S and those that did not. If the 2S chromosome segment was preferentially transmitted through both the male and female gametes, only those that carried the 2S chromosome would be viable and as a result all the progeny observed in the  $BC_3/BC_4$  self-populations would be homozygous for the 2S chromosome segment. Alternatively, if the 2S chromosome segment was not preferentially transmitted through the male or the female gametes, then the progeny would segregate for the presence of either one or two 2S chromosome segments. All progeny observed carried two copies of the 2S chromosome segment (Fig. 5), thus demonstrating that the 2S chromosome segment was preferentially transmitted through both the male and female gametes.

The frequency of transmission of chromosome segments from other linkage groups in each of the backcross progenies was much lower than that observed for linkage group 2. Of the remaining linkage groups, segments from linkage group 5 of *Ae. speltoides* had the highest frequency of transmission in all the backcross progenies observed (Table 4)

## DISCUSSION

Aegilops speltoides is a potentially important source of genetic variation for a range of agronomically important traits (Jia et al., 1996; Mago et al., 2009; Klindworth et al., 2012; Deek and Distelfeld, 2014). In the past, it has been difficult to detect and characterize wheat/wild relative introgressions due to the absence of high-throughput, genome-wide marker technologies. In this paper we have utilized an Affymetrix SNP array (King et al., 2017) complemented by a specific crossing strategy and the use of an automated, high-throughput microscope system for GISH image capture.

In this work we generated  $F_1$  interspecific hybrids between wheat and *Aegilops speltoides* and backcrossed these to wheat (Fig. 1). Hybrids were haploid for the A, B and D genomes of

TABLE 4. Transmission frequencies of the seven linkage groups of Ae. speltoides in the backcross populations  $(BC_1 \text{ to } BC_5)$  analysed by SNP genotyping

$\begin{tabular}{ c c c c c c c c c c c c c c c c c c c$							
Number of plants genotyped2225212311722536Linkage group 1 (%)18 (82)124 (49)36 (29)13 (11)3 (14)194 (36)Linkage group 2 (%)22 (100)252 (100)123 (100)117 (100)22 (100)536 (100)Linkage group 3 (%)18 (81)135 (54)27 (22)20 (17)0 (0)200 (37)Linkage group 4 (%)15 (68)89 (35)23 (19)7 (6)0 (0)134 (25)Linkage group 5 (%)19 (86)162 (64)54 (44)32 (27)6 (27)273 (51)Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)		$BC_1$	$BC_2$	BC <sub>3</sub>	$\mathrm{BC}_4$	BC <sub>5</sub>	Total
Linkage group 1 (%)18 (82)124 (49)36 (29)13 (11)3 (14)194 (36)Linkage group 2 (%)22 (100)252 (100)123 (100)117 (100)22 (100)536 (100)Linkage group 3 (%)18 (81)135 (54)27 (22)20 (17)0 (0)200 (37)Linkage group 4 (%)15 (68)89 (35)23 (19)7 (6)0 (0)134 (25)Linkage group 5 (%)19 (86)162 (64)54 (44)32 (27)6 (27)273 (51)Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)	Number of plants genotyped	22	252	123	117	22	536
Linkage group 2 (%)22 (100)252 (100)123 (100)117 (100)22 (100)536 (100)Linkage group 3 (%)18 (81)135 (54)27 (22)20 (17)0 (0)200 (37)Linkage group 4 (%)15 (68)89 (35)23 (19)7 (6)0 (0)134 (25)Linkage group 5 (%)19 (86)162 (64)54 (44)32 (27)6 (27)273 (51)Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)	Linkage group 1 (%)	18 (82)	124 (49)	36 (29)	13 (11)	3 (14)	194 (36)
Linkage group 3 (%)18 (81)135 (54)27 (22)20 (17)0 (0)200 (37)Linkage group 4 (%)15 (68)89 (35)23 (19)7 (6)0 (0)134 (25)Linkage group 5 (%)19 (86)162 (64)54 (44)32 (27)6 (27)273 (51)Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)	Linkage group 2 (%)	22 (100)	252 (100)	123 (100)	117 (100)	22 (100)	536 (100
Linkage group 4 (%)15 (68)89 (35)23 (19)7 (6)0 (0)134 (25)Linkage group 5 (%)19 (86)162 (64)54 (44)32 (27)6 (27)273 (51)Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)	Linkage group 3 (%)	18 (81)	135 (54)	27 (22)	20 (17)	0 (0)	200 (37)
Linkage group 5 (%)19 (86)162 (64)54 (44)32 (27)6 (27)273 (51)Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)	Linkage group 4 (%)	15 (68)	89 (35)	23 (19)	7 (6)	0 (0)	134 (25)
Linkage group 6 (%)17 (77)115 (46)31 (25)25 (21)0 (0)188 (35)Linkage group 7 (%)13 (59)80 (32)15 (12)9 (8)0 (0)117 (22)	Linkage group 5 (%)	19 (86)	162 (64)	54 (44)	32 (27)	6 (27)	273 (51)
Linkage group 7 (%)         13 (59)         80 (32)         15 (12)         9 (8)         0 (0)         117 (22)	Linkage group 6 (%)	17 (77)	115 (46)	31 (25)	25 (21)	0 (0)	188 (35)
	Linkage group 7 (%)	13 (59)	80 (32)	15 (12)	9 (8)	0 (0)	117 (22)

Comparative synteny of Aegilops speltoides and hexaploid wheat

FIG. 4. Synteny of *Ae. speltoides* (genetic position in cM) with hexaploid wheat (physical position in Mb) [visualized using Circos v. 0.67 (Krzywinski *et al.*, 2009)].

wheat and the S genome of *Ae. speltoides*. Thus, in the absence of homologous chromosomes the only recombination occurring at meiosis was between homoeologous chromosomes. This would normally be prevented by the presence of *Ph1*. However, the presence of the pairing promoters *Su1-Ph1* and *Su2-Ph1* located on chromosomes 3S and 7S of *Ae. speltoides* (Chen and Dvorak, 1984; Dvorak *et al.*, 2006) enabled homoeologous recombination to occur.

Since the interspecific  $F_1$  hybrids generated were haploid for the A, B, D and S genomes, their fertility was predicted to be low and this was found to be the case (Table 1). However, their fertility of 29 % was higher than that previously observed in those between wheat and *Amblyopyrum muticum*, which was 16.2 % (King *et al.*, 2017). The low fertility of the  $F_1$  hybrids resulted in the generation of only 22 BC<sub>1</sub> seeds that grew to maturity and set seed. Hence, if recombination did not occur in later generations, i.e. in the gametes of the BC<sub>1</sub>, BC<sub>2</sub>, BC<sub>3</sub> and BC<sub>4</sub> generated was limited to the 22 female  $F_1$  gametes giving rise to these 22 BC<sub>1</sub> plants. However, the level of interspecific recombination detected by the genetic mapping was such that it was possible to assemble the seven linkage groups of *Ae. speltoides*. Using the genetic linkage map of *Ae. speltoides*,



FIG. 5. GISH of a complete one-cell metaphase spread of a genotype produced by self-fertilization (BC<sub>3</sub>F<sub>3</sub>-6B) with labelled genomic *Ae. speltoides* DNA as probe, showing a homozygous pair of *Ae. speltoides* linkage group 2 introgressed chromosomes (green).

we estimated that 294 wheat/Ae. speltoides introgressions were generated spanning the entire Ae. speltoides genome. We were able to characterize these introgressions and track them through the backcross generations (Fig. 3). However, while using the genetic map to characterize the introgressions, it was important to treat the cM distances with caution, as the maps were not produced using proper mapping families. Tracking the introgressions through the different backcross generations via the SNP data showed that the majority of introgressions generated had occurred due to recombination in the  $F_1$  gametes and were therefore present in the BC, plants.

Validation of introgressions identified by SNP analysis was carried out by GISH analysis. In most cases the number of wheat/*Ae. speltoides* introgressions detected by the SNP analysis corresponded to the number of introgressions detected by the GISH analysis (Table 3 and Fig. 3). The one exception (BC<sub>5</sub>-4A) can be explained in two possible ways. The first possibility is that linkage group 2S of *Ae. speltoides* has recombined with a wheat chromosome. Although the telomere is on a separate chromosome, the SNP markers would have detected

the two introgressions as a complete chromosome. The second possibility is that the genetic linkage map of *Ae. speltoides* is not complete and therefore the smaller introgression has not been detected at all and the large segment has appeared complete in the SNP analysis.

Multicolour GISH analysis was carried out to verify which of the wheat genomes were involved in recombination with *Ae. speltoides* (Fig. 6). The majority of recombination events were shown to have occurred between *Ae. speltoides* and the B genome of wheat (91 %). Recombination had also occurred between *Ae. speltoides* and the D genome (18 %) but we found no evidence of recombination between *Ae. speltoides* and the wheat A genome. As the B genome progenitor of wheat is thought to be *Ae. speltoides* or a close relative (Sarkar and Stebbins, 1956; Riley *et al.*, 1958; Dvorak and Zhang, 1990; Maestra and Naranjo, 1998), the higher level of recombination between *Ae. speltoides* and the B genome was expected. The low level of recombination observed with the D genome of wheat would also suggest that *Ae. speltoides* is more closely related to the D genome than to the A genome. This situation



FIG. 6. Genomic *in situ* hybridization showing recombination between *Ae. speltoides* and the B and D genomes of wheat. (A) GISH analysis of a 41-chromosome  $BC_3F_1$  individual. Metaphase spread with labelled genomic *Ae. speltoides* DNA as probe showing *Ae. speltoides* (green) introgressions (white arrows). (B) Same metaphase spread as (A) with three-colour GISH showing *Ae. speltoides* recombined with the D genome (red arrows). Enlarged inset shows one of the recombined chromosomes. This individual carries 14 A chromosomes (green), 13 B chromosomes (purple), 12 D chromosomes (red) and two wheat/*Ae. speltoides* 2S recombinant chromosomes (the segment of the D genome is red and marked with the red arrow). (C) GISH analysis of a 42-chromosome  $BC_5$  individual. Metaphase spread with labelled genomic *Ae. speltoides* (green) DNA as probe showing one complete *Ae. speltoides* chromosome (blue arrow) from linkage group 2S and one wheat/*Ae. speltoides* recombined with the B genome (yellow arrow). Enlarged insert shows the recombined chromosome (multi arrise 14 A chromosome (yellow arrow). Enlarged insert shows the recombined chromosome (blue arrow) from linkage group 2S and one wheat/*Ae. speltoides* recombined with the B genome (yellow arrow). Enlarged insert shows the recombined chromosome. This individual carries 14 A chromosomes (purple), 14 D chromosomes (red), one complete *Ae. speltoides* chromosome (purple) and one wheat/*Ae. speltoides* recombined with the B genome (yellow arrow). Enlarged insert shows the recombined chromosome. This individual carries 14 A chromosomes (purple), 14 D chromosomes (red), one complete *Ae. speltoides* chromosome (purple) and one wheat/*Ae. speltoides* recombined chromosomes (purple).

closely mirrors that observed between wheat and *Amblyopyrum muticum*, where most of the recombination occurred between the chromosomes of the wild relative and the B and D genomes of wheat (King *et al.*, 2017). In addition, although the genetic location of the *Ph1* pairing suppressors located on *Am. muticum* are not currently known, the fact that both *Ae. speltoides* and *Am. muticum* carry suppressors indicates that the two species may have had a common ancestry.

We are currently unable to use the SNP markers to identify which of the chromosomes of wheat have recombined with the Ae. speltoides chromosomes. We are currently developing wheat chromosome-specific Kompetitive Allele Specific PCR (KASP) markers from the SNP markers on the 35K array. These markers will allow the analysis of large numbers of individuals, firstly to tag individual introgressions and track them through the generations (both via backcrossing and selfing) and secondly to identify which wheat chromosome(s) is (are) involved in each introgression once homozygous introgressions have been generated. Although considerably more labour-intensive and technically demanding, the latter could also be achieved using fluorescence in situ hybridization (FISH) with repetitive DNA sequences combined with GISH. This has been successfully demonstrated to identify the recipient wheat chromosomes with introgressions from Thinopyrum bessarabicum (Patokar et al., 2016), Th. intermedium and Secale cereale (Ali et al., 2016) and Aegilops markgrafii, Aegilops triuncialis and Aegilops cylindrica (Molnár et al., 2015).

In the *Ae. speltoides* accession used in this study, chromosome 2S was found to be preferentially transmitted at a frequency of 100 % through both the male and female gametes. This demonstrates that the gametocidal genes on chromosome 2S are extremely potent and resemble those in species such as *Aegilops sharonensis* (King *et al.*, 1991*a, b*; King and Laurie, 1993). The transmission of the other six linkage groups of *Ae. speltoides* was also observed in the female gametes, and although linkage group 5 showed a potentially elevated rate of transmission, there is no evidence at this time to suggest that it carries a gene for preferential transmission.

It is important to note that the work undertaken was based on a single accession of *Ae. speltoides* and thus it is not possible to determine whether all other accessions carry gametocidal genes on chromosome 2S or other linkage groups. However, if they do, then it will be necessary to remove them if introgression lines without them are to be developed. In order to develop lines that lack the 2S gametocidal genes, but still carry introgressions from other linkage groups, we are currently undertaking research to mirror the work of Friebe *et al.* (2003), who successfully deleted the genes responsible for preferential transmission in *Ae. sharonensis*.

## CONCLUSION

Researchers have been working to transfer genetic variation from its wild relatives into wheat for many years. However, although there have been some notable successes, we have barely scratched the surface of the genetic variation available for exploitation. This has been as a direct result of: (1) the difficulty of inducing wheat/wild relative introgressions and (2) an inability to detect the introgressions. The data obtained here clearly demonstrate that *Ae. speltoides* chromosomes recombine with the chromosomes of wheat at a very high frequency, giving rise to large numbers of introgressions in derivatives of the  $F_1$  hybrid. Considering the importance of the genetic variation identified in *Ae. speltoides* for a range of agronomic traits (Jia *et al.*, 1996; Mago *et al.*, 2009; Klindworth *et al.*, 2012; Deek and Distelfeld, 2014), wheat/*Ae. speltoides* introgressions have the potential to play a critical role in the development of superior wheat varieties in the future.

#### **ACKNOWLEDGEMENTS**

This work was supported by the Biotechnology and Biological Sciences Research Council as part of the Wheat Improvement Strategic Programme [grant number BB/J004596/1]. The funding body played no role in the design of the study, collection, analysis and interpretation of data and in writing the manuscript.

#### LITERATURE CITED

- Ali N, Heslop-Harrison JS, Ahmad H, Graybosch RA, Hein GL, Schwarzacher T. 2016. Introgression of chromosome segments from multiple alien species in wheat breeding lines with wheat streak mosaic virus resistance. *Heredity* 117: 114–123.
- Brozynska M, Furtado A, Henry RJ. 2015. Genomics of crop wild relatives: expanding the gene pool for crop improvement. *Plant Biotechnology Journal* 14: 1070–1085.
- Charmet G. 2011. Wheat domestication: lessons for the future. Comptes Rendus Biologies 334: 212–220.
- Chen KC, Dvorak J. 1984. The inheritance of genetic variation in *Triticum* speltoides affecting heterogenetic chromosome pairing in hybrids with *Triticum aestivum*. Canadian Journal of Genetics and Cytology 26: 279–287.
- **Deek J, Distelfeld A. 2014.** Characterisation of *Aegilops speltoides* introgression lines conferring resistance to wheat leaf rust. *Phytoparasitica* **42**: 651–672.
- Doussinault G, Delibes A, Sanchezmonge R, Garcia-Olmedo F. 1983. Transfer of a dominant gene for resistance to eyespot disease from a wild grass to hexaploid wheat. *Nature* 303: 698–700.
- Dvorak J. 1972. Genetic variability in *Aegilops speltoides* affecting homoeologous pairing in wheat. *Canadian Journal of Genetics and Cytology* 14: 371–380.
- **Dvorak J, Akhunov E. 2005.** Tempos of gene locus deletions and duplications and their relationship to recombination rate during diploid and polyploid evolution in the *Aegilops-Triticum* alliance. *Genetics* **171**: 323–332.
- Dvorak J, Zhang HB. 1990. Variation in repeated nucleotide sequences sheds light on the phylogeny of the wheat B and G genomes. *Proceedings of the National Academy of Sciences of the USA* 87: 9640–9644.
- **Dvorak J, di Terlizzi P, Zhang HB, Resta P. 1993**. The evolution of polyploid wheats identification of the A-genome donor species. *Genome* **36**: 21–31.
- Dvorak J, Deal KR, Luo M-C. 2006. Discovery and mapping of wheat Ph1 suppressors. Genetics 174: 17–27.
- Finch RA, Miller TE, Bennett MD. 1984. "Cuckoo" Aegilops addition chromosome in wheat ensures its transmission by causing chromosome breaks in meiospores lacking it. *Chromosoma* 90: 84–88.
- Friebe B, Gill BS. 1996. Chromosome banding and genome analysis in diploid and cultivated polyploid wheats. In: Jauhar PP. ed. *Methods of genome* analysis in plants. New York: CRC Press, 39–59.
- Friebe B, Zhang P, Nasuda S, Gill BS. 2003. Characterization of a knock-out mutation at the Gc2 locus in wheat. Chromosoma 111: 509–517.
- Grassini P, Eskridge KM, Cassman KG. 2013. Distinguishing between yield advances and yield plateaus in historical crop production trends. *Nature Communications* 3: 2918.
- Graybosch RA, Peterson CJ, Baenziger PS et al. 2009. Registration of 'MACE' hard red winter wheat. Journal of Plant Registration 3: 51–56.
- Haldane JBS. 1919. The probable errors of calculated linkage values, and the most accurate method of determining gametic from certain zygotic series. *Journal of Heredity* 8: 291–297.
- Haussmann BIG, Parzies HK, Presterl T, Susic Z, Miedaner T. 2004. Plant genetic resources in crop improvement. *Plant Genetic Resources* 2: 3–21.

- Jia J, Devos KM, Chao S, Miller TE, Reader SM, Gale MD. 1996. RFLPbased maps of the homoeologous group 6 chromosomes of wheat and their application in the tagging of *Pm12*, a powdery mildew resistance gene transferred from *Aegilops speltoides* to wheat. *Theoretical and Applied Genetics* 92: 559–565.
- Kato A, Lamb JC, Birchler JA. 2004. Chromosome painting using repetitive DNA sequences as probes for somatic chromosome identification in maize. *Proceedings of the National Academy of Sciences of the USA* 101: 13554–13559.
- **King IP, Laurie DA. 1993.** Chromosome damage in early embryo and endosperm development in crosses involving the preferentially transmitted 4S<sup>1</sup> chromosome of *Aegilops sharonensis. Heredity* **70**: 52–59.
- King IP, Koebner RMD, Schlegel R, Reader SM, Miller TE, Law CN. 1991a. Exploitation of a preferentially transmitted chromosome from *Aegilops sharonensis* for the elimination of segregation for height in semidwarf bread wheat varieties. *Genome* 34: 944–949.
- King IP, Miller TE, Koebner RMD. 1991b. Determination of the transmission frequency of chromosome 4S<sup>1</sup> of Aegilops sharonensis in a range of wheat genetic backgrounds. *Theoretical and Applied Genetics* 81: 519–523.
- King J, Gustafson P, Allen A, King I. 2016. Exploitation of interspecific diversity in wheat. In: Bonjean AP, Angus WJ, van Ginkel M. eds. *The* world wheat book: a history of wheat breeding, Vol. 3. Paris: Lavoisier, 1125–1139.
- King J, Grewal S, Yang C, et al. 2017. A step change in the transfer of interspecific variation into wheat from Amblyopyrum muticum. Plant Biotechnology Journal 15: 217–226.
- Klindworth DL, Niu Z, Chao S, et al. 2012. Introgressions and characterisation of a goatgrass gene for a high level of resistance to Ug99 stem rust in tetraploid wheat. Gene, Genomes and Genetics 2: 665–673.
- Krzywinski M, Schein J, Birol I, et al. 2009. Circos: an information aesthetic for comparative genomics. *Genome Research* 19: 1639–1645.
- Liu CJ, Atkinson MD, Chinoy CN, Devos KM, Gale MD. 1992. Nonhomoeologous translocations between group 4, 5 and 7 chromosomes within wheat and rye. *Theoretical and Applied Genetics* 83: 305–312.
- Maestra B, Naranjo T. 1998. Homoeologous relationships of Aegilops speltoides chromosomes to bread wheat. Theoretical and Applied Genetics 97: 181–186.
- Mago R, Zhang P, Banana HS, et al. 2009. Development of wheat lines carrying stem rust resistance gene Sr39 with reduced Aegilops speltoides chromatin and simple PCR markers for marker-assisted selection. Theoretical and Applied Genetics 119: 1441–1450.
- Maia N. 1967. Obtention de blés tendres résistants au piétin-verse (*Cercosporella herpotrichoides*) par crôisements interspécifiques blés x Aegilops. Comptes Rendus de l'Académie d'Agriculture de France 53: 149–155.
- Marcussen T, Sandve SR, Heier L, et al. 2014. Ancient hybridizations among the ancestral genomes of bread wheat. Science 345: 1250092.
- McCouch S, Baute GJ, Bradeen J, et al. 2013. Agriculture: feeding the future. Nature 499: 23–24.
- McFadden E, Sears ER. 1946. The origin of *Triticum spelta* and its free threshing hexaploid relatives. *Journal of Heredity* 37: 81–107.
- Milne I, Shaw P, Stephen G, et al. 2010. Flapjack graphical genotype visualisation. Bioinformatics 26: 3133–3134.
- Molnár I, Vrána J, Farkas A, et al. 2015. Flow sorting of C-genome chromosomes from wild relatives of wheat Aegilops markgrafii, Ae. triuncialis and Ae. cylindrica and their molecular organization. Annals of Botany 116: 189–200.

- Naranjo T, Roca A, Goicoechea PG, Giraldez R. 1987. Arm homoeology of wheat and rye chromosomes. *Genome* 29: 873–882.
- **Ogihara Y, Hasegawa K, Tsujimoto K.1994.** High-resolution cytological mapping of the long arm of chromosome 5A in common wheat using a series of deletion lines induced by gametocidal (*Gc*) genes of *Aegilops speltoides. Molecular and General Genetics* **244**: 253–259.
- van Ooijen JW. 2011. Multipoint maximum likelihood mapping in a full-sib family of outbreeding species. *Genetics Research* **93**: 343–349.
- Patokar C, Sepsi A, Schwarzacher T, Kishii M, Heslop-Harrison JS. 2016. Molecular cytogenetic characterization of novel wheat-*Thinopyrum* bessarabicum recombinant lines carrying intercalary translocations. Chromosoma 125: 163–172.
- Ray DK, Mueller ND, West PC, Foley JA. 2013. Yield trends are insufficient to double global crop production by 2050. *PLoS ONE* 8: e66428.
- Riley R, Unrau J, Chapman V. 1958. Evidence on the origin of the B genome of wheat. *Journal of Heredity* 49: 91–98
- Sarkar P, Stebbins GL.1956. Morphological evidence concerning the origin of the B genome in wheat. *American Journal of Botany* **43**: 297–304.
- Sears ER. 1955. An induced gene transfer from Aegilops to Triticum. Genetics 40: 595.
- Tanksley SD, McCouch SR. 1997. Seed banks and molecular maps: unlocking genetic potential from the wild. *Science* 277: 1063–1066.
- Tester M, Langridge P. 2010. Breeding technologies to increase crop production in a changing world. *Science* 327: 818–822.
- Tiwari VK, Wang S, Sehgal S, et al. 2014. SNP discovery for mapping alien introgressions in wheat. BMC Genomics 15: 271.
- Tiwari VK, Wang S, Danilova T, et al. 2015. Exploring the tertiary gene pool of bread wheat: sequence assembly and analysis of chromosome 5M9g0 of Aegilops geniculata. Plant Journal 84: 733–746.
- Tsujimoto H, Tsunewaki K. 1984. Gc genes in wheat and its relatives. I. Genetic analyses in common heat of a Gc gene derived from Aegilops speltoides. Canadian Journal of Genetics and Cytology 26: 78–84.
- Tsujimoto H, Tsunewaki K. 1988. Gametocidal genes in wheat and its relatives. III. Chromosome location and effects of two Aegilops speltoidesderived Gc genes in common wheat. Genome 30: 239–244.
- Tsunewaki K. 1996. Plasmon analysis as the counterpart of genome analysis. In: Jauhar PP. ed. *Methods of genome analysis in plants*. New York: CRC Press, 271–299.
- Warschefsky E, Penmetsa RV, Cook DR, von Wettberg EJB. 2014. Back to the wilds: tapping evolutionary adaptations for resilient crops through systematic hybridization with crop wild relatives. *American Journal of Botany* 101: 1791–1800.
- Wilkinson PA, Winfield MO, Barker GLA, et al. 2012. CerealsDB 2.0: an integrated resource for plant breeders and scientists. BMC Bioinformatics 13: 219.
- Wilkinson PA, Winfield MO, Barker GLA, et al. 2016. CerealsDB 3.0: expansion of resources and data integration. BMC Bioinformatics 17: 256.
- Winfield MO, Allen AM, Burridge AJ, et al. 2016. High-density SNP genotyping array for hexaploid wheat and its secondary and tertiary gene pool. *Plant Biotechnology Journal* 14: 1195–1206.
- Zhang H, Bian Y, Gou X, et al. 2013. Persistent whole-chromosome aneuploidy is generally associated with nascent allohexaploid wheat. Proceedings of the National Academy of Sciences of the USA 110: 3447–3452.
- Zhang H, Mittal N, Leamy LJ, Barazani O, Song B-H. 2017. Back into the wild – apply untapped genetic diversity of wild relatives for crop improvement. *Evolutionary Applications* 10: 5–24.